

2021

## STATISTICS — HONOURS

Fifth Paper

(Group - A)

Full Marks : 50

*The figures in the margin indicate full marks.**Candidates are required to give their answers in their own words as far as practicable.**Notation have their usual significance.*

## Unit - I

1. Attempt **any two** questions from the following : 5×2(a) Suppose  $R = (\rho_{ij})_{i,j=1(1)k}$  is the correlation matrix for a  $k$ -dimensional random vector. Show that  $|R| \leq 1$  and also interpret the case when  $|R| = 1$ .(b) If a random vector  $X = (X_1, X_2, \dots, X_p)' \sim N_p(\mu, \Sigma)$ , evaluate  $E \left( e^{\sum_{j=1}^p X_j} \right)$ .(c) Let  $X = (X_1, X_2, \dots, X_p)' \sim N_p(\mu_1, I_p)$  independently of  $Y = (Y_1, Y_2, \dots, Y_p)' \sim N_p(\mu_2, \Sigma)$ ,  $\Sigma > 0$ .Also let  $U = \sum_{i=1}^p \frac{1}{2}(X_i - \mu_{1i})$  and  $V = \sum_{i=1}^p \frac{1}{2}(X_i - \mu_{2i})$ , then prove that  $\text{Var}(U'V) = p$ .(d) If  $(N_1, N_2, \dots, N_m) \sim \text{Multinomial}(n | p_1, p_2, \dots, p_m)$ , find the condition mean and variance of  $N_i$ ,given  $\sum_{j=1}^m N_j = c$ , for  $0 \leq c < n$ .2. Attempt **any one** question from the following :(a) (i) Let  $X, Y$  and  $Z$  be independent random variables having same variance. Define the random variables  $W_1 = (X - Z)/\sqrt{2}$ ,  $W_2 = (X + Y + Z)/\sqrt{3}$  and  $W_3 = (X + 2Y + Z)/\sqrt{6}$ . Find the multiple correlation coefficient of  $W_1$  on  $(W_2, W_3)$ .(ii) Suppose  $(X_1, X_2, \dots, X_p) \sim \text{Multinomial}(n | \pi_1, \pi_2, \dots, \pi_p)$  and  $c_i$ 's are non-zero real numbers such that  $\sum_{i=1}^p c_i \pi_i = 0$ . If  $U = \sum_{i=1}^p c_i X_i$  and  $V = \sum_{i=1}^p (X_i / c_i)$ , verify whether two random variables  $U$  and  $V$  are linearly independent or not. 7+8

Please Turn Over

- (b) (i) For any non null vector  $\alpha \in \mathbb{R}'$ , establish that  $\alpha'X \sim N(\alpha'\mu, \alpha'\Omega\alpha) \Leftrightarrow X \sim N_p(\mu, \Omega)$  where  $X$  is a  $p$ -dimensional random vector and  $\Omega > 0$ .
- (ii) Let  $X_0, X_1, \dots, X_n$  be  $n+1$  independent standard normal variables. Define the random variables

$$Y_j = \rho X_0 + \sqrt{(1-\rho^2)}X_j, j=1,2,\dots,n.$$

Obtain the distribution of  $Y = (Y_1, Y_2, \dots, Y_n)'$ . Also express the partial correlation coefficient  $\rho_{12 \cdot 34 \dots n}$  in terms of  $\rho$ . 7+8

### Unit - II

3. Attempt **any two** questions from the following : 5×2

- (a) Suppose in a group of  $n$  people selected at random from a community contains  $m (< n)$  men of which  $m_0$  are smokers. Among  $(n-m)$  women in the group there are  $f_0$  smokers. Construct Pearsonian Chi-square test to judge whether a man is more prone to smoking habit than a woman.
- (b) Let  $T_n$  denote the mean of  $n$  independent random observations having common pdf  $f_\theta(x) = 2(\theta-x)/\theta^2, 0 < x < \theta, \theta > 0$ .

Applying delta method, show that  $\sqrt{n}(\log T_n - \log(\theta/3)) \xrightarrow{D} N(0, 1/2)$ , as  $n \rightarrow \infty$ .

- (c) Based on  $n$  independent copies of  $X \sim N(\mu, \sigma^2)$ , obtain the large sample standard error of sample coefficient of variation.
- (d) Find the asymptotic distribution of sample median of a random sample of size  $n$  drawn from the distribution with pdf  $f_\theta(x) = 2\theta^2/x^3, x > \theta, \theta > 0$ .

4. Attempt **any one** question from the following :

- (a) (i) Explain the term 'convergence in distribution'. If, as  $n \rightarrow \infty, X_n \xrightarrow{D} X$  and  $Y_n \xrightarrow{P} Y$ , prove that  $X_n + Y_n \xrightarrow{D} X + Y$ .

- (ii) State the central limit theorem for a sequence of iid random variables. For a sequence  $\{X_k\}_{k \geq 1}$  of independent Poisson ( $\lambda$ ) variables prove that, as  $n \rightarrow \infty$ ,

$$\sqrt{n}(\sqrt{S_n/n} - \lambda\sqrt{n/S_n}) \xrightarrow{D} N(0,1)$$

where  $S_n = X_1 + X_2 + \dots + X_n$ . (2+6)+(2+5)

- (b) (i) Based on two independent random samples of sizes  $n_1$  and  $n_2$  from  $N(\mu_1, \sigma_1^2)$  and  $N(\mu_2, \sigma_2^2)$  respectively, describe a large sample test procedure for testing the null hypothesis  $H_0: \sigma_1 = \sigma_2$ . If  $H_0$  is rejected, then determine  $100(1-\alpha)\%$  confidence interval for  $|\sigma_1 - \sigma_2|$ .
- (ii) Suppose  $(X_1, X_2, \dots, X_n)$  represents a random sample drawn from  $N(\mu, \sigma^2)$  population. Derive the asymptotic distribution of sample  $r$ -th order central moment. (5+3)+7